

Assignment #2

Due on Friday September 14, 2007

Read Chapter 2 on *Vector Algebra*, pp. 29–49, in Bressoud.

Do the following problems

1. Recall that the dot product, or inner product, of two vectors in \mathbb{R}^n is symmetric, bi-linear and positive definite; that is, for vectors v, v_1, v_2 and w in \mathbb{R}^n ,

(i) $v \cdot w = w \cdot v$

(ii) $(c_1v_1 + c_2v_2) \cdot w = c_1v_1 \cdot w + c_2v_2 \cdot w$, and

(iii) $v \cdot v \geq 0$ for all $v \in \mathbb{R}^n$ and $v \cdot v = 0$ if and only if v is the zero vector.

Use these properties of the the inner product in \mathbb{R}^n to derive the following properties of the norm $\|\cdot\|$ in \mathbb{R}^n , where

$$\|v\| = \sqrt{v \cdot v} \quad \text{for all vectors } v \in \mathbb{R}^n.$$

(a) $\|v\| \geq 0$ for all $v \in \mathbb{R}^n$ and $\|v\| = 0$ if and only if $v = \vec{0}$.

(b) For a scalar c , $\|cv\| = |c|\|v\|$.

2. Recall the Cauchy-Schwarz inequality: For any vectors v and w in \mathbb{R}^n ,

$$|v \cdot w| \leq \|v\|\|w\|.$$

Use this inequality to derive the triangle inequality: For any vectors v and w in \mathbb{R}^n ,

$$\|v + w\| \leq \|v\| + \|w\|.$$

(*Suggestion:* Start with the expression $\|v + w\|^2$ and use the properties of the inner product to simplify it.)

3. Given two non-zero vectors v and w in \mathbb{R}^n , the cosine of the angle, θ , between the vectors can be defined by

$$\cos \theta = \frac{v \cdot w}{\|v\|\|w\|}.$$

Use the Cauchy-Schwarz inequality to justify why this definition makes sense.

4. Two vectors v and w in \mathbb{R}^n are said to be *orthogonal* or perpendicular, if and only if $v \cdot w = 0$.

Show that if v and w are orthogonal, then

$$\|v + w\|^2 = \|v\|^2 + \|w\|^2.$$

Give a geometric interpretation of this result in two-dimensional Euclidean space.

5. A vector u in \mathbb{R}^n is said to be a unit vector if and only if $\|u\| = 1$. Let u be a unit vector in \mathbb{R}^n and v be any vector in \mathbb{R}^n .

- (a) Give the parametric equation of the line through origin in the direction of u .
- (b) Let $f(t) = \|v - tu\|^2$ for all $t \in \mathbb{R}$. Explain why this function gives the square of the distance from the point at v to a point on the line through the origin in the direction of u .
- (c) Show that $f(t)$ is minimized when $t = v \cdot u$.