

## Assignment #7

Due on Monday, October 12, 2009

Read Section 7.1 on *Limits*, pp. 171–178, in Bressoud.

## Background and Definitions

- (*Open Set*) A subset,  $U$ , of  $\mathbb{R}^n$  is said to be **open** if for any  $x \in U$  there exists a positive number  $r$  such that  $B_r(x) = \{y \in \mathbb{R}^n \mid \|y - x\| < r\}$  is entirely contained in  $U$ . (The empty set,  $\emptyset$ , is considered to be an open set.)
- (*Continuous Functions 1*) Let  $U$  denote an open subset of  $\mathbb{R}^n$ . A function  $F: U \rightarrow \mathbb{R}^m$  is said to be continuous at  $x \in U$  if and only if

$$\lim_{\|y-x\| \rightarrow 0} \|F(y) - F(x)\| = 0.$$

If  $F$  is continuous at every point in  $U$ , then  $F$  is said to be continuous **on**  $U$ .

- (*Pre-image*) If  $B \subseteq \mathbb{R}^m$ , the *pre-image* of  $B$  under the map  $F: U \rightarrow \mathbb{R}^m$ , denoted by  $F^{-1}(B)$ , is defined as the set  $F^{-1}(B) = \{x \in U \mid F(x) \in B\}$ .  
Note that  $F^{-1}(B)$  is always defined even if  $F$  does not have an inverse map.
- (*Continuous Functions 2*) Let  $U$  denote an open subset of  $\mathbb{R}^n$ . A function  $F: U \rightarrow \mathbb{R}^m$  is continuous on  $U$  if and only if, for every open subset  $V$  of  $\mathbb{R}^m$ , the pre-image of  $V$  under  $F$ ,  $F^{-1}(V)$  is open in  $\mathbb{R}^n$ .
- (*Composition of Continuous Functions*) Let  $U$  denote an open subset of  $\mathbb{R}^n$  and  $Q$  an open subset of  $\mathbb{R}^m$ . Suppose that the maps  $F: U \rightarrow \mathbb{R}^m$  and  $G: Q \rightarrow \mathbb{R}^k$  are continuous on their respective domains and that  $F(U) \subseteq Q$ . Then, the composition  $G \circ F: U \rightarrow \mathbb{R}^k$  is continuous on  $U$ .

Do the following problems

1. Let  $U$  denote an open subset of  $\mathbb{R}^n$ . Suppose that  $f: U \rightarrow \mathbb{R}$  is a scalar field and  $G: U \rightarrow \mathbb{R}^m$  is vector valued function.
  - (a) Explain how the product  $fG$  is defined.
  - (b) Prove that if both  $f$  and  $G$  are continuous on  $U$ , then the vector valued function  $fG$  is also continuous on  $U$ .

2. Let  $U$  be an open subset of  $\mathbb{R}^2$ . Let  $f: U \rightarrow \mathbb{R}$  and  $g: U \rightarrow \mathbb{R}$  be two scalar fields on  $U$ , and define  $h: U \rightarrow \mathbb{R}$  by

$$h(x, y) = f(x, y)g(x, y) \quad \text{for all } (x, y) \in U.$$

Prove that if both  $f$  and  $g$  are continuous on  $U$ , then so is  $h$ .

*Suggestion:* First prove that the function  $G: \mathbb{R}^2 \rightarrow \mathbb{R}$ , defined by  $G(x, y) = xy$  for all  $(x, y) \in \mathbb{R}^2$ , is continuous. Then, let  $F: U \rightarrow \mathbb{R}^2$  denote the map given by

$$F(x, y) = (f(x, y), g(x, y)) \quad \text{for all } (x, y) \in U,$$

and observe that

$$h = G \circ F.$$

3. Let  $U = \mathbb{R}^n \setminus \{\mathbf{0}\} = \{v \in \mathbb{R}^n \mid v \neq \mathbf{0}\}$ .

- (a) Prove that  $U$  is an open subset of  $\mathbb{R}^n$ .  
(b) Define  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  by

$$f(v) = \frac{1}{\|v\|} \quad \text{for all } v \in U.$$

Prove that  $f$  is continuous on  $U$ .

*Suggestion:* Note that the function,  $g$ , defined by

$$g(t) = \frac{1}{t} \quad \text{for all } t \neq 0,$$

is continuous for  $t \neq 0$ .

4. Let  $I \subseteq \mathbb{R}$  be an open interval and  $\sigma: I \rightarrow \mathbb{R}^n$  be continuous path in  $\mathbb{R}^n$  satisfying  $\sigma(t) \neq \mathbf{0}$  for all  $t \in I$ . Define the function  $f: I \rightarrow \mathbb{R}$  by

$$f(t) = \frac{1}{\|\sigma(t)\|} \quad \text{for all } t \in I.$$

Prove that  $f$  is continuous on  $I$ .

5. Exercise 12 on page 180 in the text.