

## Solutions to Review Problems for Exam 1

1. There are 5 red chips and 3 blue chips in a bowl. The red chips are numbered 1, 2, 3, 4, 5 respectively, and the blue chips are numbered 1, 2, 3 respectively. If two chips are to be drawn at random and without replacement, find the probability that these chips are have either the same number or the same color.

**Solution:** Let  $R$  denote the event that the two chips are red. Then the assumption that the chips are drawn at random and without replacement implies that

$$\Pr(R) = \frac{\binom{5}{2}}{\binom{8}{2}} = \frac{5}{14}.$$

Similarly, if  $B$  denotes the event that both chips are blue, then

$$\Pr(B) = \frac{\binom{3}{2}}{\binom{8}{2}} = \frac{3}{28}.$$

It then follows that the probability that both chips are of the same color is

$$\Pr(R \cup B) = \Pr(R) + \Pr(B) = \frac{13}{28},$$

since  $R$  and  $B$  are disjoint.

Let  $N$  denote the event that both chips show the same number. Then,

$$\Pr(N) = \frac{3}{\binom{8}{2}} = \frac{3}{28}.$$

Finally, since  $R \cup B$  and  $N$  are disjoint, then the probability that the chips are have either the same number or the same color is

$$\Pr(R \cup B \cup N) = \Pr(R \cup B) + \Pr(N) = \frac{13}{28} + \frac{3}{28} = \frac{16}{28} = \frac{4}{7}.$$

□

2. A person has purchased 10 of 1,000 tickets sold in a certain raffle. To determine the five prize winners, 5 tickets are drawn at random and without replacement. Compute the probability that this person will win at least one prize.

**Solution:** Let  $N$  denote the event that the person will not win any prize. Then

$$\Pr(N) = \frac{\binom{995}{10}}{\binom{1000}{10}}; \quad (1)$$

that is, the probability of purchasing 10 non-winning tickets.

It follows from (1) that

$$\begin{aligned} \Pr(N) &= \frac{(990)(989)(988)(987)(986)}{(1000)(999)(998)(997)(996)} \\ &= \frac{435841667261}{458349513900} \\ &\approx 0.9509. \end{aligned} \quad (2)$$

Thus, using the result in (2), the probability of the person winning at least one of the prizes is

$$\begin{aligned} \Pr(N^c) &= 1 - \Pr(N) \\ &\approx 1 - 0.9509 \\ &= 0.0491, \end{aligned}$$

or about 4.91%. □

3. Let  $(\mathcal{C}, \mathcal{B}, \Pr)$  denote a probability space, and let  $E_1$ ,  $E_2$  and  $E_3$  be mutually disjoint events in  $\mathcal{B}$ . Find  $\Pr[(E_1 \cup E_2) \cap E_3]$  and  $\Pr(E_1^c \cup E_2^c)$ .

**Solution:** Since  $E_1$ ,  $E_2$  and  $E_3$  are mutually disjoint events, it follows that  $(E_1 \cup E_2) \cap E_3 = \emptyset$ ; so that

$$\Pr[(E_1 \cup E_2) \cap E_3] = 0.$$

Next, use De Morgan's law to compute

$$\begin{aligned} \Pr(E_1^c \cup E_2^c) &= \Pr([E_1 \cap E_2]^c) \\ &= \Pr(\emptyset^c) \\ &= \Pr(\mathcal{C}) \\ &= 1. \end{aligned}$$

□

4. Let  $(\mathcal{C}, \mathcal{B}, \Pr)$  denote a probability space, and let  $A$  and  $B$  events in  $\mathcal{B}$ . Show that

$$\Pr(A \cap B) \leq \Pr(A) \leq \Pr(A \cup B) \leq \Pr(A) + \Pr(B). \quad (3)$$

**Solution:** Since  $A \cap B \subseteq A$ , it follows that

$$\Pr(A \cap B) \leq \Pr(A). \quad (4)$$

Similarly, since  $A \subseteq A \cup B$ , we get that

$$\Pr(A) \leq \Pr(A \cup B). \quad (5)$$

Next, use the identity

$$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B),$$

and fact that that

$$\Pr(A \cap B) \geq 0,$$

to obtain that

$$\Pr(A \cup B) \leq \Pr(A) + \Pr(B). \quad (6)$$

Finally, combine (4), (5) and (6) to obtain (3). □

5. Let  $(\mathcal{C}, \mathcal{B}, \Pr)$  denote a probability space, and let  $E_1$ ,  $E_2$  and  $E_3$  be mutually independent events in  $\mathcal{B}$  with probabilities  $\frac{1}{2}$ ,  $\frac{1}{3}$  and  $\frac{1}{4}$ , respectively. Compute the exact value of  $\Pr(E_1 \cup E_2 \cup E_3)$ .

**Solution:** First, use De Morgan's law to compute

$$\Pr[(E_1 \cup E_2 \cup E_3)^c] = \Pr(E_1^c \cap E_2^c \cap E_3^c) \quad (7)$$

Then, since  $E_1$ ,  $E_2$  and  $E_3$  are mutually independent events, it follows from (7) that

$$\Pr[(E_1 \cup E_2 \cup E_3)^c] = \Pr(E_1^c) \cdot \Pr(E_2^c) \cdot \Pr(E_3^c),$$

so that

$$\begin{aligned} \Pr[(E_1 \cup E_2 \cup E_3)^c] &= (1 - \Pr(E_1))(1 - \Pr(E_2))(1 - \Pr(E_3)) \\ &= \left(1 - \frac{1}{2}\right) \left(1 - \frac{1}{3}\right) \left(1 - \frac{1}{4}\right) \\ &= \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{3}{4}, \end{aligned}$$

so that

$$\Pr[(E_1 \cup E_2 \cup E_3)^c] = \frac{1}{4}. \quad (8)$$

It then follows from (8) that

$$\Pr(E_1 \cup E_2 \cup E_3) = 1 - \Pr[(E_1 \cup E_2 \cup E_3)^c] = \frac{3}{4}.$$

□

6. Let  $(\mathcal{C}, \mathcal{B}, \Pr)$  denote a probability space, and let  $E_1$ ,  $E_2$  and  $E_3$  be mutually independent events in  $\mathcal{B}$  with  $\Pr(E_1) = \Pr(E_2) = \Pr(E_3) = \frac{1}{4}$ . Compute  $\Pr[(E_1^c \cap E_2^c) \cup E_3]$ .

**Solution:** First, use De Morgan's law to compute

$$\Pr[((E_1^c \cap E_2^c) \cup E_3)^c] = \Pr[(E_1^c \cap E_2^c)^c \cap E_3^c] \quad (9)$$

Next, use the assumption that  $E_1$ ,  $E_2$  and  $E_3$  are mutually independent events to obtain from (9) that

$$\Pr[((E_1^c \cap E_2^c) \cup E_3)^c] = \Pr[(E_1^c \cap E_2^c)^c] \cdot \Pr[E_3^c], \quad (10)$$

where

$$\Pr[E_3^c] = 1 - \Pr(E_3) = \frac{3}{4}, \quad (11)$$

and

$$\begin{aligned} \Pr[(E_1^c \cap E_2^c)^c] &= 1 - \Pr[E_1^c \cap E_2^c] \\ &= 1 - \Pr[E_1^c] \cdot \Pr[E_2^c], \end{aligned} \quad (12)$$

by the independence of  $E_1$  and  $E_2$ .

It follows from the calculations in (13) that

$$\begin{aligned}
 \Pr[(E_1^c \cap E_2^c)^c] &= 1 - (1 - \Pr[E_1])(1 - \Pr[E_2]) \\
 &= 1 - \left(1 - \frac{1}{4}\right) \left(1 - \frac{1}{4}\right) \\
 &= 1 - \frac{3}{4} \cdot \frac{3}{4} \\
 &= \frac{7}{16}
 \end{aligned} \tag{13}$$

Substitute (11) and the result of the calculations in (13) into (10) to obtain

$$\Pr[((E_1^c \cap E_2^c) \cup E_3)^c] = \frac{7}{16} \cdot \frac{3}{4} = \frac{21}{64}. \tag{14}$$

Finally, use the result in (14) to compute

$$\begin{aligned}
 \Pr[(E_1^c \cap E_2^c) \cup E_3^c] &= 1 - \Pr[((E_1^c \cap E_2^c) \cup E_3)^c] \\
 &= 1 - \frac{21}{64} \\
 &= \frac{43}{64}.
 \end{aligned}$$

□

7. A bowl contains 5 chips of the same size and shape. One the chips is red and the rest are blue. Draw chips from the bowl at random, one at a time and without replacement, until the red chip is drawn.

(a) Describe the sample space of this experiment.

**Solution:** Denoting the red chip by  $R$  and any of the blue chips by  $B$ , we have that the sample space for this experiment is

$$\mathcal{C} = \{R, BR, BBR, BBBR, BBBBR\}.$$

□

(b) Define the probability function for this experiment. Justify your answer.

**Solution:** Since we are assuming that the chips are drawn at random and without replacement, we have that

$$\Pr(R) = \frac{1}{5};$$

$$\Pr(BR) = \frac{4}{5} \cdot \frac{1}{4} = \frac{1}{5};$$

$$\Pr(BBR) = \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{1}{3} = \frac{1}{5};$$

$$\Pr(BBBR) = \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{2}{3} \cdot \frac{1}{2} = \frac{1}{5};$$

and

$$\Pr(BBBBBR) = \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{2}{3} \cdot \frac{1}{2} \cdot \frac{1}{1} = \frac{1}{5}.$$

Thus, we conclude that

$$\Pr(c) = \frac{1}{5}, \quad \text{for all } c \in \mathcal{C}.$$

□

(c) Compute the probability that at least two draws will be needed to get the red chip.

**Solution:** The event,  $E$ , that at least two draws will be needed to get the red chip, is the complement of the set  $\{R\}$ . Thus,  $E = \{R\}^c$  and therefore

$$\Pr(E) = 1 - \Pr(\{R\}) = 1 - \frac{1}{5} = \frac{4}{5}.$$

□

8. Dreamboat cars are produced at three different factories A, B and C. Factory A produces 20 percent of the total output of Dreamboats, B produces 50 percent, and C produces 30 percent. However, 5 percent of the cars produced at A are lemons, 2 percent of those produced at B are lemons, and 10 percent of those produced at C are lemons. If you buy a Dreamboat and it turns out to be lemon, what is the probability that it was produced at factory A?

**Solution:** Let  $A$  denote the event that the car was produced in Factory A,  $B$  the event the car was made in Factory B, and  $C$  the event the car was made in Factory C. We then have that

$$\Pr(A) = 0.20, \quad \Pr(B) = 0.50 \quad \text{and} \quad \Pr(C) = 0.30.$$

Let  $L$  denote the event that a given car is a lemon. We are then given the conditional probabilities

$$\Pr(L | A) = 0.05, \quad \Pr(L | B) = 0.02, \quad \text{and} \quad \Pr(L | C) = 0.10.$$

We want to compute  $\Pr(A | L)$ ,

$$\Pr(A | L) = \frac{\Pr(A \cap L)}{\Pr(L)},$$

where

$$\Pr(A \cap L) = \Pr(A) \cdot \Pr(L | A) = (0.20) \cdot (0.05) = 0.01,$$

and

$$\begin{aligned} \Pr(L) &= \Pr(A) \cdot \Pr(L | A) + \Pr(B) \cdot \Pr(L | B) + \Pr(C) \cdot \Pr(L | C) \\ &= (0.20) \cdot (0.05) + (0.50) \cdot (0.02) + (0.30) \cdot (0.10) \\ &= 0.01 + 0.01 + 0.03 \\ &= 0.05. \end{aligned}$$

Hence,

$$\Pr(A | L) = \frac{0.01}{0.05} = \frac{1}{5},$$

or 20%. □

9. Let  $(\mathcal{C}, \mathcal{B}, \Pr)$  denote a probability space, and let  $A$  and  $B$  events in  $\mathcal{B}$ . Given that  $\Pr(A) = 1/3$ ,  $\Pr(B) = 1/5$  and  $\Pr(A | B) + \Pr(B | A) = 2/3$ , compute  $\Pr(A^c \cup B^c)$ .

**Solution:** Assume that

$$\Pr(A) = \frac{1}{3}, \quad \Pr(B) = \frac{1}{5}, \tag{15}$$

and

$$\Pr(A | B) + \Pr(B | A) = \frac{2}{3}. \tag{16}$$

First, use De Morgan's Law and the Rule of Complements to compute

$$\begin{aligned} \Pr(A^c \cup B^c) &= \Pr((A \cap B)^c) \\ &= 1 - \Pr(A \cap B); \end{aligned}$$

so that

$$\Pr(A^c \cup B^c) = 1 - \Pr(A) \cdot \Pr(B | A). \quad (17)$$

Thus, we need to compute  $\Pr(B | A)$ . To do so, first use

$$\Pr(A | B) = \frac{\Pr(A \cap B)}{\Pr(B)}$$

to obtain

$$\Pr(A | B) = \frac{\Pr(A) \cdot \Pr(B | A)}{\Pr(B)},$$

or

$$\Pr(A | B) = \frac{5}{3} \cdot \Pr(B | A), \quad (18)$$

in view of (15). Next, combine (18) and (16) to obtain

$$\frac{5}{3} \cdot \Pr(B | A) + \Pr(B | A) = \frac{2}{3},$$

from which we get

$$\Pr(B | A) = \frac{1}{4}.$$

Using this value in (17) and the value of  $\Pr(A)$  in (15) we obtain that

$$\Pr(A^c \cup B^c) = 1 - \frac{1}{3} \cdot \frac{1}{4},$$

from which we get that

$$\Pr(A^c \cup B^c) = \frac{11}{12}.$$

□

10. Let  $(\mathcal{C}, \mathcal{B}, \Pr)$  denote a probability space, and let  $A$  and  $B$  independent events in  $\mathcal{B}$  with  $\Pr(B) > 0$ . Given that  $\Pr(A) = 1/3$ , compute  $\Pr(A \cup B^c | B)$ .

**Solution:** Use the definition of conditional probability to compute

$$\Pr(A \cup B^c | B) = \frac{\Pr((A \cup B^c) \cap B)}{\Pr(B)}, \quad (19)$$

where, by the distributive property,

$$(A \cup B^c) \cap B = (A \cap B) \cup (B^c \cap B) = (A \cap B) \cup \emptyset = A \cap B;$$

so that,

$$\Pr((A \cup B^c) \cap B) = \Pr(A \cap B),$$

and, using the assumption of independence of  $A$  and  $B$ ,

$$\Pr((A \cup B^c) \cap B) = \Pr(A) \cdot \Pr(B).$$

Consequently, in view of (19),

$$\Pr(A \cup B^c \mid B) = \Pr(A) = \frac{1}{3}.$$

□