

Assignment #4

Due on Friday, September 29, 2017

Read Section 4.1, *Gâteaux Differentiability*, in the class lecture notes at <http://pages.pomona.edu/~ajr04747/>

Background and Definitions

Gâteaux Differentiability. Let V denote a normed linear space, and V_o a non-trivial subspace of V . Let $J: V \rightarrow \mathbb{R}$ be a functional defined on V . We say that J is Gâteaux differentiable at $u \in V$ in the direction of $v \in V_o$ if

$$\left. \frac{d}{dt} J(u + tv) \right|_{t=0} = \lim_{t \rightarrow 0} \frac{J(u + tv) - J(u)}{t} \text{ exists.}$$

If the limit exists, we denote it by $dJ(u; v)$, and call it the Gâteaux derivative of J at u in the direction of v , or the first variation of J at u in the direction of v .

Do the following problems

1. Let $V = C^1([a, b], \mathbb{R})$ and $V_o = C_o^1([a, b], \mathbb{R})$. Define

$$J(y) = \frac{1}{2} \int_a^b (y'(x))^2 dx, \quad \text{for all } y \in C^1([a, b], \mathbb{R}).$$

Show that $J: V \rightarrow \mathbb{R}$ is Gâteaux differentiable at every $y \in V$ in the direction of $v \in V_o$, and compute $dJ(y; v)$ for all $y \in V$ and $v \in V_o$.

2. Let $G: \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function and $J: C^1([a, b], \mathbb{R}) \rightarrow \mathbb{R}$ be the functional given by

$$J(y) = \int_a^b x^2 (y'(x))^3 dx + G(y(b)) \quad \text{for all } y \in C^1([a, b], \mathbb{R}).$$

Show that J is Gâteaux differentiable at every $y \in C^1([a, b], \mathbb{R})$ in the direction of $\eta \in C_o^1([a, b], \mathbb{R})$ and compute $dJ(y; \eta)$ for all $y \in C^1([a, b], \mathbb{R})$ and $\eta \in C_o^1([a, b], \mathbb{R})$.

3. Let V be a normed linear space and $L: V \rightarrow \mathbb{R}$ be a linear function. Prove that L is Gâteaux differentiable in V in the direction of any $v \in V$ and compute $dL(u; v)$ for any $u \in V$ and $v \in V$.

4. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ denote a C^1 real valued function of a single variable, and define $J: C^1([a, b], \mathbb{R}) \rightarrow \mathbb{R}$ by $J(y) = \int_a^b f(y'(x)) \, dx$ for all $y \in C^1([a, b], \mathbb{R})$. Show that J is Gâteaux differentiable at every $y \in C^1([a, b], \mathbb{R})$ in the direction of every $\eta \in C_o^1([a, b], \mathbb{R})$, and compute the Gâteaux derivative of J at every $y \in C^1([a, b], \mathbb{R})$ in the direction of every $\eta \in C_o^1([a, b], \mathbb{R})$.
5. Let V denote a normed linear space and V_o a non-trivial subspace of V . Assume that $J: V \rightarrow \mathbb{R}$ is Gâteaux differentiable at every $u \in V$ in the direction of every $v \in V_o$. Suppose that J has a local minimum at $u \in V$; so that,

$$J(u) \leq J(w), \quad \text{for all } w \in V \text{ with } \|w - u\| < \delta,$$

and some $\delta > 0$. Show that

$$dJ(u; v) = 0, \quad \text{for all } v \in V_o$$