

## Assignment #12

Due on Monday, November 5, 2018

**Read** Section 3.6 on *The Chain Rule and the Rate of Change along a Path*, pp. 133–136, in Baxandall and Liebek’s text.

**Read** Section 3.7 on *Directional Derivatives*, pp. 138–141, in Baxandall and Liebek’s text.

**Read** Section 3.8 on *The Gradient and Smooth Surfaces*, pp. 142–151, in Baxandall and Liebek’s text.

**Read** Section 4.6 on *Derivatives of Compositions* in the class Lecture Notes at <http://pages.pomona.edu/~ajr04747/>.

**Do** the following problems

- Let  $U = \{(x, y) \in \mathbb{R}^2 \mid x \neq 0\}$  and define  $f: U \rightarrow \mathbb{R}$  by  $f(x, y) = \arctan\left(\frac{y}{x}\right)$ , for all  $(x, y) \in U$ .
  - Compute the gradient of  $f$  in  $U$ .
  - Let  $I$  be an open interval and  $\sigma: I \rightarrow U$  be a differentiable path given by  $\sigma(t) = (x(t), y(t))$  for  $t \in I$ . Define  $\theta: I \rightarrow \mathbb{R}$  by  $\theta(t) = (f \circ \sigma)(t)$  for all  $t \in I$ . Apply the Chain Rule to verify that  $\theta' = \frac{-yx' + xy'}{x^2 + y^2}$ .
  - Apply the result from part (b) to the path  $\sigma: \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}^2$  given by  $\sigma(t) = (\cos t, \sin t)$ , for  $t \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ .
- Suppose that the temperature in a region of space is given by a function  $T: \mathbb{R}^3 \rightarrow \mathbb{R}$  given by

$$T(x, y, z) = kx^2(y - z), \quad \text{for all } (x, y, z) \in \mathbb{R}^3,$$

and some positive constant  $k$ .

An insect flies in the region along a path modeled by a  $C^1$  function  $\sigma: \mathbb{R} \rightarrow \mathbb{R}^3$ . Suppose that at time  $t = 0$  the insect is located at  $(0, 0, 0)$  and its velocity is  $\sigma'(0) = \hat{i} + \hat{j} + 2\hat{k}$ . Compute the rate of change of temperature sensed by the insect at time  $t = 0$ .

3. Let  $I$  be an open interval of real numbers and  $U$  be an open subset of  $\mathbb{R}^n$ . Suppose that  $\sigma: I \rightarrow \mathbb{R}^n$  is a differentiable path and that  $f: U \rightarrow \mathbb{R}$  is a differentiable scalar field. Assume also that the image of  $I$  under  $\sigma$ ,  $\sigma(I)$ , is contained in  $U$ . Suppose also that the derivative of the path  $\sigma$  satisfies

$$\sigma'(t) = -\nabla f(\sigma(t)) \quad \text{for all } t \in I.$$

Show that if the gradient of  $f$  along the path  $\sigma$  is never zero, then  $f$  decreases along the path as  $t$  increases.

*Suggestion:* Use the Chain Rule to compute the derivative of  $f(\sigma(t))$ .

4. A set  $U \subseteq \mathbb{R}^n$  is said to be **path connected** iff for any vectors  $x$  and  $y$  in  $U$ , there exists a differentiable path  $\sigma: [0, 1] \rightarrow \mathbb{R}^n$  such that  $\sigma(0) = x$ ,  $\sigma(1) = y$  and  $\sigma(t) \in U$  for all  $t \in [0, 1]$ ; i.e., any two elements in  $U$  can be connected by a differentiable path whose image is entirely contained in  $U$ .

Suppose that  $U$  is an open, path connected subset of  $\mathbb{R}^n$ . Let  $f: U \rightarrow \mathbb{R}$  be a differentiable scalar field such that  $\nabla f(x)$  is the zero vector for all  $x \in U$ . Prove that  $f$  must be constant.

5. Let  $U$  be an open subset of  $\mathbb{R}^n$  and  $f: U \rightarrow \mathbb{R}$  be a differentiable scalar field defined on  $U$ . The function  $f$  is said to be homogeneous of order  $k$  if

$$f(tv) = t^k f(v),$$

for all  $v \in U$  and all positive  $t \in \mathbb{R}$  such that  $tv \in U$ .

- Show that the function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  given by  $f(x, y) = xy$ , for all  $(x, y) \in \mathbb{R}^2$ , is homogeneous of order 2.
- Give examples of a scalar field which is homogeneous of order 1 and of a scalar field which is homogeneous of order 0.
- Prove Euler's Theorem: If  $f: U \rightarrow \mathbb{R}$  is differentiable and homogeneous of order  $k$ , then

$$x_1 \frac{\partial f}{\partial x_1}(x) + x_2 \frac{\partial f}{\partial x_2}(x) + \cdots + x_n \frac{\partial f}{\partial x_n}(x) = kf(x),$$

for all  $x = (x_1, x_2, \dots, x_n) \in U$ .

*Suggestion:* Let  $\sigma: \mathbb{R} \rightarrow \mathbb{R}^n$  be given by  $\sigma(t) = tx$ , for all  $t \in \mathbb{R}$  and  $x = (x_1, x_2, \dots, x_n) \in U$ , and apply the Chain Rule to the composition  $f \circ \sigma$ .