

Assignment #4

Due on Friday, September 27, 2019

Read Section 2.5 on *The Cross Product* in the class Lecture Notes at <http://pages.pomona.edu/~ajr04747/>.

Do the following problems

1. Let u , v and w denote non-zero vectors in \mathbb{R}^3 . Given that $u \cdot w = 0$, $u \cdot v = c$, where c is a real constant, and $u \times v = w$, find the components of v in each of the three mutually orthogonal directions: u , w and $u \times w$.
2. Prove that the cross product is non-associative; that is, find three vectors u , v and w in \mathbb{R}^3 such that $(u \times v) \times w \neq u \times (v \times w)$.
3. Let v and w denote vectors in \mathbb{R}^3 , and $\mathbf{0}$ the zero-vector in \mathbb{R}^3 .
 - (a) Prove that if $v \times w = \mathbf{0}$ and $v \cdot w = 0$, then at least one of v or w must be the zero vector.
 - (b) Prove that $v \cdot (v \times w) = 0$.

4. In this problem and the next, we derive the vector identity

$$u \times (v \times w) = (u \cdot w)v - (u \cdot v)w$$

for any vectors u , v and w in \mathbb{R}^3 .

- (a) Argue that $u \times (v \times w)$ lies in the span of v and w . Consequently, there exist scalars t and s such that

$$u \times (v \times w) = tv + sw$$

- (b) Show that $(u \cdot v)t + (u \cdot w)s = 0$.

5. Let u , v and w be as in the previous problem.

- (a) Use the results of the previous problem to conclude that there exists a scalar r such that

$$u \times (v \times w) = r[(u \cdot w)v - (u \cdot v)w].$$

- (b) By considering some simple examples, deduce that $r = 1$ in the previous identity