

Review Problems for Exam 1

1. Consider the set $B = \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}$.
- (a) Show that B is a basis for \mathbb{R}^2 .
- (b) Give the coordinates of the vector $v = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ relative to B . Interpret your result geometrically.
2. Give a basis for the span of the following set of vectors in \mathbb{R}^4

$$\left\{ \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} -2 \\ 0 \\ 3 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -3 \\ 6 \\ -3 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ -4 \\ 1 \end{pmatrix} \right\}.$$

3. Find a basis for the solution space of the system

$$\begin{cases} x_1 - x_2 + x_3 - x_4 = 0 \\ 2x_1 - x_2 - 2x_4 = 0 \\ -x_1 + x_3 + x_4 = 0, \end{cases}$$

and compute its dimension.

4. Prove that any set of four vectors in \mathbb{R}^3 must be linearly dependent.
5. Show that if the set $\{v_1, v_2\}$ is a linearly independent subset of \mathbb{R}^n , then so is the set $\{v_1, cv_1 + v_2\}$, where c is a scalar, and, conversely, if $\{v_1, cv_1 + v_2\}$ is linearly independent, then so is $\{v_1, v_2\}$. Show also that $\text{span}\{v_1, v_2\} = \text{span}\{v_1, cv_1 + v_2\}$.
6. Let J and H be planes in \mathbb{R}^3 given by

$$J = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \mid 2x + 3y - 6z = 0 \right\} \quad \text{and} \quad H = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \mid x - 2y + z = 0 \right\}.$$

- (a) Give bases for J and H and compute their dimensions.
- (b) Give a basis for the subspace $J \cap H$ and compute $\dim(J \cap H)$.

7. Let W be a subspace of \mathbb{R}^n .
- (a) Prove that if $v \in W$ and $v \neq \mathbf{0}$, then $rv = sv$ implies that $r = s$, where r and s are scalars.
 - (b) Prove that if W has more than one element, then W has infinitely many elements.
8. Let W be a subspace of \mathbb{R}^n and S_1 and S_2 be subsets of W .
- (a) Show that $\text{span}(S_1 \cap S_2) \subseteq \text{span}(S_1) \cap \text{span}(S_2)$.
 - (b) Give an example in which $\text{span}(S_1 \cap S_2) \neq \text{span}(S_1) \cap \text{span}(S_2)$.
 - (c) Show that if $S_1 \subseteq S_2$ and S_2 is linearly independent, then S_1 is also linearly independent.
 - (d) Show that if $S_1 \subseteq S_2$ and S_1 is linearly dependent, then S_2 is also linearly dependent.
9. Let W_1 and W_2 be two subspaces of \mathbb{R}^n . We write $W_1 \oplus W_2$ for the subspace $W_1 + W_2$ for the special case in which $V = W_1 \cap W_2 = \{\mathbf{0}\}$. Show that every vector $v \in W_1 \oplus W_2$ can be written in the form $v = v_1 + v_2$, where $v_1 \in W_1$ and $v_2 \in W_2$, in one and only one way; that is, if $v = u_1 + u_2$, where $u_1 \in W_1$ and $u_2 \in W_2$, then $u_1 = v_1$ and $u_2 = v_2$.
10. Let $v \in \mathbb{R}^n$ and define $W = \{w \in \mathbb{R}^n \mid \langle w, v \rangle = 0\}$.
- (a) Prove that W is a subspace of \mathbb{R}^n .
 - (b) Suppose that $v \neq \mathbf{0}$ and compute $\dim(W)$.