

## Assignment #4

Due on Wednesday, February 5, 2014

**Read** Sections 2.4 on *Defining a Probability Function* in the class lecture notes at <http://pages.pomona.edu/~ajr04747/>

**Read** Section 1.5 on *The Definition of Probability* in DeGroot and Schervish.

**Read** Section 1.6 on *Finite Sample Spaces* in DeGroot and Schervish.

**Do** the following problems

1. Let  $(\mathcal{C}, \mathcal{B}, \Pr)$  be a sample space. Suppose that  $E_1, E_2, E_3, \dots$  is a sequence of events in  $\mathcal{B}$  satisfying

$$E_1 \supseteq E_2 \supseteq E_3 \supseteq \dots$$

Prove that  $\lim_{n \rightarrow \infty} \Pr(E_n) = \Pr\left(\bigcap_{k=1}^{\infty} E_k\right)$ .

*Hint:* Use the analogous result for an increasing nested sequence of events presented in class and De Morgan's laws.

2. A point  $(x, y)$  is to be selected at random from a square  $S$  containing all the points  $(x, y)$  such that  $0 \leq x \leq 1$  and  $0 \leq y \leq 1$ . Suppose that the probability that the selected point will belong to each specified subset of  $S$  is equal to the area of that subset. Find the probability of each of the following subsets:

(a) the subset of points such that  $\left(x - \frac{1}{2}\right)^2 + \left(y - \frac{1}{2}\right)^2 \geq \frac{1}{4}$ ;

(b) the subset of points such that  $\frac{1}{2} < x + y < \frac{3}{2}$ ;

(c) the subset of points such that  $y < 1 - x^2$ ;

(d) the subset of points such that  $x = y$ .

3. In a random experiment, two balanced dice are rolled.
  - (a) What is the probability that the sum of the two numbers that appear will be even?
  - (b) What is the probability that the difference of the two numbers that appear will be less than 3?

4. A coin is tossed as many times as necessary to turn up one head. Thus, the elements of the sample space  $\mathcal{C}$  corresponding to this experiment are

$$H, TH, TTH, TTTH, \dots$$

Let  $\Pr$  be a function that assigns to these elements the values  $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$  respectively.

- (a) Show that  $\Pr(\mathcal{C}) = 1$ .
- (b) Let  $E_1$  denote the event  $E_1 = \{H, TH, TTH, TTTH \text{ or } TTTTH\}$ , and compute  $\Pr(E_1)$ .
- (c) Let  $E_2 = \{TTTTH, TTTTTH\}$ , and compute  $\Pr(E_2)$ ,  $\Pr(E_1 \cap E_2)$  and  $\Pr(E_2 \setminus E_1)$
5. Let  $\mathcal{C} = \{x \in \mathbb{R} \mid x > 0\}$  and define  $\Pr$  on open intervals  $(a, b)$  with  $0 < a < b$  by

$$\Pr((a, b)) = \int_a^b e^{-x} \, dx.$$

- (a) Show that  $\Pr(\mathcal{C}) = 1$ .
- (b) Let  $E = \{x \in \mathcal{C} \mid 4 < x < \infty\}$ , and compute  $\Pr(E)$ ,  $\Pr(E^c)$  and  $\Pr(E \cup E^c)$ .