

Assignment #21

Due on Friday, November 30, 2007

Read on *The Fundamental Theorem of Calculus*, pp. 279–320 in Chapter 10 of Bressoud’s book.

Background and Definitions

Green’s Theorem. The Fundamental Theorem of Calculus,

$$\int_M d\omega = \int_{\partial M} \omega,$$

takes the following form in two–dimensional Euclidean space:

Let R denote a region in \mathbb{R}^2 bounded by a simple closed curve, ∂R , made up of a finite number of C^1 paths traversed in the counterclockwise sense. Let P and Q denote two C^1 scalar fields defined on some open set containing R and its boundary, ∂R . Then,

$$\int_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy = \int_{\partial R} P dx + Q dy. \quad (1)$$

Do the following problems

1. Apply Green’s Theorem, as expressed in the formula (1), to the functions $P(x, y) = -y$ and $Q(x, y) = x$ to derive the formula

$$\text{area}(R) = \frac{1}{2} \int_{\partial R} -y dx + x dy. \quad (2)$$

to compute the area of the region R .

2. Use the formula (2) derived in the previous theorem to compute the area enclosed by the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1,$$

where a and b are positive real numbers.

3. In Problem 2(b) of Assignment #20, you showed that another form of Fundamental Theorem of Calculus is

$$\int_R \operatorname{div}(F) \, dx dy = \text{Flux of } F \text{ across } \partial R;$$

that is, the flux of F across the boundary of R is the double integral of the divergence of F over the region R . Writing $F = P \hat{i} + Q \hat{j}$, the above form of the Fundamental Theorem of Calculus takes the form

$$\int_R \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} \right) dx dy = \int_{\partial R} F \cdot \hat{n} \, ds, \quad (3)$$

where \hat{n} is a unit vector perpendicular to ∂R and pointing to the outside of ∂R .

Use formula (3) to compute the flux of the field $F = x \hat{i} + y \hat{j}$ across the square with vertices $(-1, -1)$, $(1, -1)$, $(1, 1)$ and $(-1, 1)$.

4. Let f and g be two scalar fields defined on some open subset of \mathbb{R}^2 . Suppose that f and g are C^1 and that ∇g is a C^1 vector field. Show that

$$\operatorname{div}(f \nabla g) = \nabla f \cdot \nabla g + f \operatorname{div}(\nabla g).$$

$\operatorname{div}(\nabla g)$ is called the *Laplacian* of g and is usually denoted by Δg ; thus,

$$\Delta g = \frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2}.$$

5. Let f and g be as in the previous problem. Use formula (3) and the result from the previous problem to show that

$$\int_R f \Delta g \, dx dy = \int_{\partial R} f \frac{\partial g}{\partial n} \, ds - \int_R \nabla f \cdot \nabla g \, dx dy,$$

where $\frac{\partial g}{\partial n}$ denotes the derivative of g in the direction of \hat{n} , or $D_{\hat{n}}g$.