

## Assignment #6

Due on Friday September 28, 2007

Read Section 7.1 on *Limits*, pp. 171–178, in Bressoud.

## Background and Definitions

- Let  $U$  denote an open subset of  $\mathbb{R}^n$ . A function  $F: U \rightarrow \mathbb{R}^m$  is said to be continuous at  $x \in U$  if and only if

$$\lim_{\|y-x\| \rightarrow 0} \|F(y) - F(x)\| = 0.$$

- If  $A \subseteq U$ , the *image* of  $A$  under the map  $F: U \rightarrow \mathbb{R}^m$ , denoted by  $F(A)$ , is defined as the set

$$F(A) = \{y \in \mathbb{R}^m \mid y = F(x) \text{ for some } x \in A\}.$$

- If  $B \subseteq \mathbb{R}^m$ , the *pre-image* of  $B$  under the map  $F: U \rightarrow \mathbb{R}^m$ , denoted by  $F^{-1}(B)$ , is defined as the set

$$F^{-1}(B) = \{x \in U \mid F(x) \in B\}.$$

Note that  $F^{-1}(B)$  is always defined even if  $F$  does not have an inverse map.

**Do** the following problems

1. Use the triangle inequality to prove that, for any  $x$  and  $y$  in  $\mathbb{R}^n$ ,

$$\left| \|y\| - \|x\| \right| \leq \|y - x\|.$$

Use this inequality to deduce that the function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  given by

$$f(x) = \|x\| \quad \text{for all } x \in \mathbb{R}^n$$

is continuous on  $\mathbb{R}^n$ .

2. Let  $f(x, y)$  and  $g(x, y)$  denote two functions defined on a open region,  $D$ , in  $\mathbb{R}^2$ . Prove that the vector field  $F: D \rightarrow \mathbb{R}^2$ , defined by

$$F \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} f(x, y) \\ g(x, y) \end{pmatrix} \quad \text{for all } \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2,$$

is continuous on  $D$  if and only if  $f$  and  $g$  are both continuous on  $D$ .

3. Let  $U$  denote an open subset of  $\mathbb{R}^n$  and let  $F: U \rightarrow \mathbb{R}^m$  and  $G: U \rightarrow \mathbb{R}^m$  be two given functions.

- (a) Explain how the sum  $F + G$  is defined.
- (b) Prove that if both  $F$  and  $G$  are continuous on  $U$ , then their sum is also continuous.  
(*Suggestion:* The triangle inequality might come in handy.)

4. In each of the following, given the function  $F: U \rightarrow \mathbb{R}^m$  and the set  $B$ , compute the pre-image  $F^{-1}(B)$ .

- (a)  $F: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,  $F \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x^2 + y^2 \\ x^2 - y^2 \end{pmatrix}$ , and  $B = \left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}$ .
- (b)  $f: D' \rightarrow \mathbb{R}$ ,

$$f(x, y) = \frac{1}{\sqrt{1 - x^2 - y^2}}, \quad \text{for } (x, y) \in D'$$

where  $D' = \{(x, y) \in \mathbb{R}^2 \mid 0 < x^2 + y^2 < 1\}$  (the punctured unit disc),  
 $B = \{1\}$ .

- (c)  $f: D' \rightarrow \mathbb{R}$  is as in part (b), and  $B = \{2\}$ .
- (d)  $f: D' \rightarrow \mathbb{R}$  is as in part (b), and  $B = \{1/2\}$ .

5. Compute the image the given sets under the following maps

- (a)  $\sigma: \mathbb{R} \rightarrow \mathbb{R}^2$ ,  $\sigma(t) = (\cos t, \sin t)$  for all  $t \in \mathbb{R}$ . Compute  $\sigma(\mathbb{R})$ .
- (b)  $f: D' \rightarrow \mathbb{R}$  and  $D'$  are as given in part (b) of the previous problem. Compute  $f(D')$ .