

## Assignment #16

Due on Friday, November 7, 2008

Read Section 3.1 on *The Calculus of Curves*, pp. 53–65, in Bressoud.

Read Section 5.2 on *Line Integrals*, pp. 113–119, in Bressoud.

Do the following problems

1. Consider a portion of a helix,  $C$ , parametrized by the path

$$\sigma(t) = (\cos t, t, \sin t) \quad \text{for } 0 \leq t \leq \pi.$$

Let  $F(x, y, z) = x \hat{i} + y \hat{j} + z \hat{k}$ , for all  $(x, y, z) \in \mathbb{R}^3$ , be a vector field in  $\mathbb{R}^3$ . Evaluate the line integral  $\int_C F \cdot T$ ; that is, the integral of the tangential component of the field  $F$  along the curve  $C$ .

2. Evaluate

$$\int_C yz \, dx + xz \, dy + xy \, dz$$

where  $C$  is the directed line segment from the point  $(1, 1, 0)$  to the point  $(3, 2, 1)$  in  $\mathbb{R}^3$ .

3. Exercises 1(a)(b)(c) on page 119 in the text.
4. Exercises 1(d)(e)(f) on page 119 in the text.
5. Let  $f: U \rightarrow \mathbb{R}$  be a  $C^1$  scalar field defined on an open subset  $U$  of  $\mathbb{R}^n$ . Define the vector field  $F: U \rightarrow \mathbb{R}^n$  by  $F(x) = \nabla f(x)$  for all  $x \in U$ . Suppose that  $C$  is a  $C^1$  simple curve in  $U$  connecting the point  $x$  to the point  $y$  in  $U$ . Show that

$$\int_C F \cdot T = f(y) - f(x).$$

Conclude therefore that the line integral of  $F$  along a path from  $x$  to  $y$  in  $U$  is independent of the path connecting  $x$  to  $y$ . The field  $F$  is called a *gradient field*.