

Assignment #12

Due on Monday, October 27, 2014

Read Section 2.12 on *Euclidean Inner Product and Norm* in the class lecture notes at <http://pages.pomona.edu/~ajr04747/>

Background and Definitions

- (*Transpose of a vector*). Given a vector $v = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$ in \mathbb{R}^n , the **transpose** of v , denoted by v^T , is the row vector

$$v^T = (x_1 \ x_2 \ \cdots \ x_n).$$

- (*Row-Column Product*). Given a row-vector, R , of dimension n and a column-vector, C , also of dimension n , we define the product RC as follows:

Write $R = [x_1 \ x_2 \ \cdots \ x_n]$ and $C = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}$; then,

$$RC = [x_1 \ x_2 \ \cdots \ x_n] \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = x_1y_1 + x_2y_2 + \cdots + x_ny_n.$$

- (*Euclidean inner product*). Given vectors $v = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$ and $w = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}$ in \mathbb{R}^n ,

the *Euclidean inner product* of v and w , denoted by $\langle v, w \rangle$, is the real number (or scalar) obtained by follows

$$\langle v, w \rangle = v^T w = (x_1 \ x_2 \ \cdots \ x_n) \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = x_1y_1 + x_2y_2 + \cdots + x_ny_n.$$

- (*Orthogonality*). Two vectors v and w in \mathbb{R}^n are said to be **orthogonal** if $\langle v, w \rangle = 0$.

- (*Euclidean norm*). Given a vector $v = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$ in \mathbb{R}^n , its **Euclidean norm**, denoted by $\|v\|$, is defined by

$$\|v\| = \sqrt{\langle v, v \rangle} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}.$$

- (*Unit vectors in \mathbb{R}^n*). A vector $u \in \mathbb{R}^n$ is said to be a **unit vector** if $\|u\| = 1$.

Do the following problems

1. The vectors $v_1 = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$, and $v_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$ span a two-dimensional subspace in \mathbb{R}^3 ; in other words, a plane through the origin. Give two unit vectors which are orthogonal to each other, and which also span the plane.
2. Let $W = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \in \mathbb{R}^3 \mid 3x - 2y + z = 0 \right\}$. Find a non-zero vector v in \mathbb{R}^3 which is orthogonal to every vector in W ; that is, $v \neq \mathbf{0}$ and

$$\langle v, w \rangle = 0 \quad \text{for all } w \in W.$$

3. Let u_1, u_2, \dots, u_n be unit vectors in \mathbb{R}^n which are mutually orthogonal; that is,

$$\langle u_i, u_j \rangle = 0 \quad \text{for } i \neq j.$$

Prove that the set $\{u_1, u_2, \dots, u_n\}$ is a basis for \mathbb{R}^n , and that, for any $v \in \mathbb{R}^n$,

$$v = \sum_{i=1}^n \langle v, u_i \rangle u_i.$$

4. The Euclidean inner product of two vectors in \mathbb{R}^n is symmetric, bi-linear and positive definite; that is, for vectors v , v_1 , v_2 and w in \mathbb{R}^n ,

- (i) $\langle v, w \rangle = \langle w, v \rangle$,
- (ii) $\langle c_1 v_1 + c_2 v_2, w \rangle = c_1 \langle v_1, w \rangle + c_2 \langle v_2, w \rangle$, and
- (iii) $\langle v, v \rangle \geq 0$ for all $v \in \mathbb{R}^n$ and $\langle v, v \rangle = 0$ if and only if v is the zero vector.

Use these properties of the the inner product in \mathbb{R}^n to derive the following properties of the norm $\| \cdot \|$ in \mathbb{R}^n :

- (a) $\|v\| \geq 0$ for all $v \in \mathbb{R}^n$ and $\|v\| = 0$ if and only if $v = \mathbf{0}$.
- (b) For a scalar c , $\|cv\| = |c|\|v\|$.

5. The Cauchy-Schwarz inequality for any vectors v and w in \mathbb{R}^n states that

$$|\langle v, w \rangle| \leq \|v\| \|w\|.$$

Use this inequality to derive the triangle inequality: For any vectors v and w in \mathbb{R}^n ,

$$\|v + w\| \leq \|v\| + \|w\|.$$

(*Suggestion:* Start with the expression $\|v + w\|^2$ and use the properties of the inner product to simplify it.)