

Assignment #16

Due on Monday, November 19, 2018

Read Section 5.4 on *The Fundamental Theorem of Calculus*, pp. 292–295, in Baxandall and Liebek’s text.

Read Section 5.5 on *Potential Functions and Conservative Fields*, pp. 296–308, in Baxandall and Liebek’s text.

Read Section 5.2 on *Line Integrals* in the class Lecture Notes at <http://pages.pomona.edu/~ajr04747/>.

Read Section 5.3 on *Gradient Fields* in the class Lecture Notes at <http://pages.pomona.edu/~ajr04747/>.

Read Section 5.4 on *Flux Across Plane Curves* in the class Lecture Notes at <http://pages.pomona.edu/~ajr04747/>.

Background and Definitions

- (*Path Connected Sets*) A set $U \subseteq \mathbb{R}^n$ is said to be path connected if and only if for any vectors p and q in U , there exists a C^1 path $\sigma: [0, 1] \rightarrow \mathbb{R}^n$ such that $\sigma(0) = p$, $\sigma(1) = q$ and $\sigma(t) \in U$ for all $t \in [0, 1]$; i.e., any two elements in U can be connected by a C^1 path whose image is entirely contained in U .
- (*Flux Across a Simple, Closed Curve in \mathbb{R}^2*) Let U denote an open subset of \mathbb{R}^2 and $F: U \rightarrow \mathbb{R}^2$ be a two-dimensional vector field given by

$$F(x, y) = P(x, y) \hat{i} + Q(x, y) \hat{j}, \quad \text{for all } (x, y) \in U,$$

where P and Q are scalar fields defined in U . Let C denote a simple, piece-wise C^1 , closed curve contained in U , which is oriented in the counterclockwise sense.

The flux of F across C , denoted by $\oint_C F \cdot \hat{n} \, ds$, is defined by

$$\oint_C F \cdot \hat{n} \, ds = \int_C P(x, y) \, dy - Q(x, y) \, dx,$$

where \hat{n} denotes the outward unit normal to the curve C , wherever it is defined.

Do the following problems

1. Integrate the 1-form $yz \, dx + xz \, dy + xy \, dz$ over each of the following curves in \mathbb{R}^3 that connect $(0, 1, 0)$ to $(2, 1, 1)$:
 - (a) the straight line from $(0, 1, 0)$ to $(2, 1, 1)$,
 - (b) the lines from $(0, 1, 0)$ to $(0, 1, 1)$ to $(2, 1, 1)$,
 - (c) the lines from $(0, 1, 0)$ to $(2, 1, 0)$ to $(2, 1, 1)$,
 - (d) the arc $(2t, (2t - 1)^2, t)$, for $0 \leq t \leq 1$.

2. Let U denote an open subset of \mathbb{R}^n that is path connected, and let $F: U \rightarrow \mathbb{R}^n$ be a vector field with the property that

$$\int_C F \cdot d\vec{r} = 0,$$

for any simple, piece-wise C^1 , closed curve, C , contained in U .

Let p and q be points in U . Since U is path connected, there exists a C^1 path, $\sigma: [0, 1] \rightarrow U$, connecting p to q . Assume that σ parametrizes a curve C_1 in U . Prove that if $\gamma: [0, 1] \rightarrow U$ is another C^1 path that connects p to q , and $C_2 = \gamma([0, 1])$ is parametrized by γ , then

$$\int_{C_1} F \cdot d\vec{r} = \int_{C_2} F \cdot d\vec{r}.$$

3. Let U denote an open subset of \mathbb{R}^n and let $F: U \rightarrow \mathbb{R}^n$ be a vector field with the property that $F(v) = \nabla f(v)$ for all $v \in U$, where $f: U \rightarrow \mathbb{R}$ is a C^1 scalar field.

Prove that if C is any C^1 , simple, closed curve in U , then

$$\int_C F \cdot d\vec{r} = 0.$$

4. Let $F(x, y) = x^2 \hat{i} + y^2 \hat{j}$ and C be the boundary of the square with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$ and $(0, 1)$, oriented in the counterclockwise sense. Compute the flux of F across C .

5. Compute the flux, $\oint_C F \cdot \hat{n} \, ds$, where $F(x, y) = x \hat{i} + y \hat{j}$, for all $(x, y) \in \mathbb{R}^2$ and C is the unit circle oriented in the counterclockwise sense.