

Assignment #19

Due on Wednesday, December 5, 2018

Read Section 11.3 on *Differential 2-Forms*, pp. 527–534, in Baxandall and Liebek's text.

Read Section 5.6 on *Calculus of Differential Forms* in the class Lecture Notes at <http://pages.pomona.edu/~ajr04747/>.

Read Section 5.7 on *Evaluating Differential 2-Forms: Double Integrals* in the class Lecture Notes at <http://pages.pomona.edu/~ajr04747/>.

Background and Definitions

- (*The Fundamental Theorem of Calculus in \mathbb{R}^2 or \mathbb{R}^3 for Oriented Triangles*) Let U denote an open region in \mathbb{R}^2 or \mathbb{R}^3 and T an oriented triangle contained in U . Denote the boundary of T by ∂T . If ω is any differential 1-form defined in U , the

$$\int_T d\omega = \oint_{\partial T} \omega. \quad (1)$$

- (*Green's Theorem for Oriented Triangles*) Let U denote an open region in \mathbb{R}^2 and T an oriented triangle contained in U . Denote the boundary of T by ∂T , and assume that it is oriented in the counterclockwise sense. For any C^1 functions, $P: U \rightarrow \mathbb{R}$ and $Q: U \rightarrow \mathbb{R}$, defined in U ,

$$\iint_T \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy = \oint_{\partial T} P dx + Q dy \quad (2)$$

- (*Divergence of a Vector Field in \mathbb{R}^2*) Given a C^1 vector field, $F(x, y) = P(x, y)\hat{i} + Q(x, y)\hat{j}$, defined on some open subset U of \mathbb{R}^2 , the divergence of F is the scalar field on U given by

$$\operatorname{div} F(x, y) = \frac{\partial P}{\partial x}(x, y) + \frac{\partial Q}{\partial y}(x, y) \quad \text{for all } (x, y) \in U. \quad (3)$$

Do the following problems

1. Evaluate the differential form $3 dx \wedge dy$ on each of the following oriented triangles:

$$(a) [(5, 2), (1, 3), (3, 4)] \quad (b) [(1, 0, -2), (3, 1, 5), (-2, 1, 0)]$$

2. Let P and Q denote smooth scalar fields defined in some open region, U , or \mathbb{R}^2 , and define the 1-form $\omega = P dy - Q dx$.

(a) Compute the differential, $d\omega$, of ω .

- (b) Recall that the integral $\int_C \omega$, where C is a simple closed curve in U , gives the flux of the field $F = P \hat{i} + Q \hat{j}$, across the curve C .

What does the Fundamental Theorem of Calculus in (1), where T is a positively oriented triangle in U , say about the flux of F across the boundary of T and the divergence of F as defined in (3)?

3. Verify the Fundamental Theorem of Calculus in (1) for the differential 1-form

$$\omega = yz dx + xz dy + xy dz,$$

and the oriented triangle $T = [P_o P_1 P_2]$, where P_o , P_1 and P_2 are any three non-collinear points in \mathbb{R}^3 .

4. Let T denote the triangle with vertices $P_o(0, 0)$, $P_1(2, 0)$ and $P_2(1, 1)$, where the boundary, ∂T , of T is oriented in the counterclockwise sense. Let $F: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the vector field given by

$$F(x, y) = -\frac{y}{2} \hat{i} + \frac{x}{2} \hat{j}.$$

Evaluate the line integral $\oint_{\partial T} F \cdot d\mathbf{r}$ by applying Green's Theorem in (2).

5. Let T and F be as in Problem 4.

Evaluate the flux of F across ∂T , $\oint_{\partial T} F \cdot d\mathbf{n}$, by applying Green's Theorem in (2).