

## Assignment #2

Due on Friday, September 20, 2019

Read Chapter 3, on *Indirect Methods*, in the class lecture notes at <http://pages.pomona.edu/~ajr04747/>

## Background and Definitions

- **Continuity.** A function  $f: [a, b] \rightarrow \mathbb{R}$  is said to be continuous at  $x_o \in [a, b]$  if for any  $\varepsilon > 0$ , there exists  $\delta > 0$  (which depends on  $\varepsilon$  and  $x_o$ ) such that  $|f(x) - f(x_o)| < \varepsilon$  for all  $x \in [a, b]$  with  $|x - x_o| < \delta$ .
- **The Class  $C([a, b], \mathbb{R})$ .** If  $f$  is continuous at every point in  $[a, b]$ , we say that  $f$  is continuous on  $[a, b]$  and write  $f \in C([a, b], \mathbb{R})$ .
- **The Class  $C_o([a, b], \mathbb{R})$ .** If  $f$  is continuous at every point in  $[a, b]$  and  $f(a) = 0$  and  $f(b) = 0$ , we write  $f \in C_o([a, b], \mathbb{R})$ .
- **The Class  $C^1([a, b], \mathbb{R})$ .** If  $f$  is differentiable in an open interval that contains  $[a, b]$ , and  $f'$  is continuous on  $[a, b]$ , we write  $f \in C^1([a, b], \mathbb{R})$ .
- **The Class  $C_o^1([a, b], \mathbb{R})$ .** If  $f \in C^1([a, b], \mathbb{R})$  and  $f(a) = f(b) = 0$ , we write  $f \in C_o^1([a, b], \mathbb{R})$ .

Do the following problems

1. Prove that if  $f \in C([a, b], \mathbb{R})$  and  $f(x_o) \neq 0$  for some  $x_o \in (a, b)$ , then there exists an interval  $(x_o - \delta, x_o + \delta)$  contained in  $(a, b)$  such that  $f(x) \neq 0$  for all  $x \in (x_o - \delta, x_o + \delta)$ .
2. Assume that  $f \in C([a, b], \mathbb{R})$  and that  $f(x) \geq 0$  for all  $x \in [a, b]$ . Prove that, if

$$\int_a^b f(x) dx = 0,$$

then  $f(x) = 0$  for all  $x \in [a, b]$ .

3. Assume that  $f \in C([a, b], \mathbb{R})$ . Suppose that

$$\int_c^d f(x) dx = 0,$$

for all  $c$  and  $d$  such that  $a \leq c < d \leq b$ . Show that  $f(x) = 0$  for all  $x \in [a, b]$ .

4. **The Fundamental Lemma in the Calculus of Variations.** Let  $f \in C([a, b], \mathbb{R})$  and suppose that

$$\int_a^b f(x)\eta(x) dx = 0, \quad \text{for all } \eta \in C_o([a, b], \mathbb{R}).$$

Show that  $f(x) = 0$  for all  $x \in [a, b]$ .

5. **The Second Fundamental Lemma in the Calculus of Variations.** In this problem we prove the second fundamental lemma in the Calculus of Variations: Let  $f \in C([a, b], \mathbb{R})$  and suppose that

$$\int_a^b f(x)\eta'(x) dx = 0, \quad \text{for all } \eta \in C_o^1([a, b], \mathbb{R}).$$

Then,  $f$  must be constant on  $[a, b]$ .

- (a) Put

$$c = \frac{1}{b-a} \int_a^b f(x) dx$$

and define  $\eta: [a, b] \rightarrow \mathbb{R}$  by

$$\eta(x) = \int_a^x (f(t) - c) dt, \quad \text{for } x \in [a, b].$$

Verify that  $\eta \in C_o^1([a, b], \mathbb{R})$ .

- (b) Show that

$$\int_a^b (f(x) - c)^2 dx = 0.$$

- (c) Deduce that  $f(x) = c$  for all  $x \in [a, b]$ .