

Assignment #3

Due on Monday, January 31, 2011

Read Section 1.2 on *The Vector Space* \mathbb{R}^n in Baxandall and Liebek's text (pp. 2–9).

Read Sections 2.3 on *The Dot Product and Euclidean Norm* in the class Lecture Notes (pp. 11–12).

Read Sections 2.4 on *Orthogonality and Projections* in the class Lecture Notes (pp. 12–16).

Do the following problems

1. The vectors $v_1 = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$ and $\vec{v}_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$ span a two-dimensional subspace in \mathbb{R}^3 , in other words, a plane through the origin. Give two unit vectors which are orthogonal to each other, and which also span the plane.

2. Use an appropriate orthogonal projection to compute the shortest distance from the point $P(1, 1, 2)$ to the plane in \mathbb{R}^3 whose equation is

$$2x + 3y - z = 6.$$

3. The dual space of \mathbb{R}^n , denoted $(\mathbb{R}^n)^*$, is the vector space of all linear transformations from \mathbb{R}^n to \mathbb{R} .

For a given $w \in \mathbb{R}^n$, define $T_w: \mathbb{R}^n \rightarrow \mathbb{R}$ by

$$T_w(v) = w \cdot v \quad \text{for all } v \in \mathbb{R}^n.$$

Show that T_w is an element of the dual of \mathbb{R}^n for all $w \in \mathbb{R}^n$.

4. Prove that for every linear transformation, $T: \mathbb{R}^n \rightarrow \mathbb{R}$, there exists $w \in \mathbb{R}^n$ such that

$$T(v) = w \cdot v \quad \text{for every } v \in \mathbb{R}^n.$$

(*Hint:* See where T takes the standard basis $\{e_1, e_2, \dots, e_n\}$ in \mathbb{R}^n .)

5. Let u_1, u_2, \dots, u_n be unit vectors in \mathbb{R}^n which are mutually orthogonal; that is,

$$u_i \cdot u_j = 0 \quad \text{for } i \neq j.$$

Prove that the set $\{u_1, u_2, \dots, u_n\}$ is a basis for \mathbb{R}^n , and that, for any $v \in \mathbb{R}^n$,

$$v = \sum_{i=1}^n (v \cdot u_i) u_i.$$