

## Assignment #3

Due on Monday, February 11, 2013

Read Section 2.3.1 on *Nondimensionalization*, in the class lecture notes at <http://pages.pomona.edu/~ajr04747/>.

Do the following problems

1. Nondimensionalize the Logistic growth equation for bacterial growth in a medium with carrying capacity  $K$ :  $\frac{dN}{dt} = rN \left(1 - \frac{N}{K}\right)$ , where  $r$  is the intrinsic growth rate of the population, by introducing dimensionless variables

$$u = \frac{N}{\mu} \quad \text{and} \quad \tau = \frac{t}{\lambda}.$$

Give interpretations for the scaling parameters  $\mu$  and  $\lambda$ .

2. Consider again the chemostat model without flow in or out of a single chamber depicted in Figure 1. Proceed as in Problems 1–4 in Assignment 1 assuming this




Figure 1: One-Compartment Chemostat Model

time that the *per capita* growth rate is given by the Michaelis-Menten enzyme kinetics relation

$$K(c) = \frac{rc}{a+c},$$

where  $c = Q/V$  is the nutrient concentration in the growth medium, to derive a differential equations for the bacterial density,  $n = N/V$ , and the nutrient concentration. You will need to use the yield  $Y = 1/\alpha$ , or the number of new cells produced in the chemostat due to consumption of one unit of nutrient.

Give an interpretation for the parameter  $r$ .

3. The differential equation

$$\frac{dN}{dt} = rN \left( 1 - \frac{N}{K} \right) - p(N, \Lambda), \quad (1)$$

models a population that is subject to predation reflected in the term  $p(N, \Lambda)$ , which depends on the population size,  $N$ , and a set of parameters,  $\Lambda$ . In the absence of predation the population undergoes logistic growth with intrinsic growth rate,  $r$ , and carrying capacity,  $K$ .

In 1978, Ludwig, Jones and Holling published an article in the Journal of Animal Ecology (*Qualitative analysis of insect outbreak systems: the spruce budworm and forest*, Volume 47, pp. 315–332) in which they proposed the following constitutive equation for the predation term,

$$p(N, a, b) = \frac{bN^2}{a^2 + N^2}. \quad (2)$$

- (a) Give interpretations for the parameters  $a$  and  $b$  in (2).  
 (b) Nondimensionalize the differential equation in (1) by introducing dimensionless variables

$$u = \frac{N}{\mu} \quad \text{and} \quad \tau = \frac{t}{\lambda},$$

to obtain the dimensionless equation

$$\frac{du}{d\tau} = \alpha u \left( 1 - \frac{u}{\beta} \right) - \frac{u^2}{1 + u^2}, \quad (3)$$

where  $\alpha$  and  $\beta$  are dimensionless parameters.

Express  $\alpha$  and  $\beta$  in terms of the parameters  $r$ ,  $K$ ,  $a$  and  $b$ .

4. Observe that  $u = 0$  is an equilibrium point of the equation in (3). Determine the nature of the stability of this equilibrium point.  
 5. The equation

$$\alpha \left( 1 - \frac{u}{\beta} \right) - \frac{u}{1 + u^2} = 0, \quad (4)$$

which yields the non-zero equilibrium points of (3), cannot be easily solved algebraically.

- (a) Explain why the equation in (4) must have at least one real solution, and at most three distinct real solutions.  
 (b) Determine conditions on  $\alpha$  and  $\beta$  that will guarantee that the equation in (4) will have (i) exactly one real solution, (ii) two distinct real solutions, and (iii) three distinct real solutions.