

## Review Problems for Exam 1

1. **Modeling the Spread of a Disease.** In a simple model for a disease that is spread through infections transmitted between individuals in a population, the population is divided into three compartments pictured in Figure 1. The




Figure 1: SIR Compartments

first compartment,  $S(t)$ , denotes the set of individuals in a population that are susceptible to acquiring the disease; the second compartment,  $I(t)$ , denotes the set of infected individual who can also infect others; and the third compartment,  $R(t)$ , denotes the set of individuals who had the disease and who have recovered from it; they can no longer get infected.

Assume that the total number of individuals in the population,

$$N = S(t) + I(t) + R(t),$$

is constant. Susceptible individuals can get infected by contact with infectious individuals and move to the infected class. This is indicated by the arrow going from the  $S(t)$  compartment to the  $I(t)$  compartment.

The rate at which susceptible individuals get infected is proportional to product of number of susceptible individuals and the number of infected individuals with constant of proportionality  $\beta > 0$ . The rate at which infected individuals recover is proportional to the number of infected individuals with constant of proportionality  $\gamma > 0$ . What are the units for  $\beta$  and  $\gamma$ ?

Use conservation principles to derive a system of differential equations for the functions  $S$ ,  $I$  and  $R$ , assuming that they are differentiable. Models of this type were first studied by Kermack and McKendrick in the early 1930s.

Introduce dimensionless variables

$$\widehat{s}(t) = \frac{S(t)}{N}, \quad \widehat{i}(t) = \frac{I(t)}{N}, \quad \widehat{r}(t) = \frac{R(t)}{N}, \quad \text{and} \quad \widehat{t} = \frac{t}{\tau},$$

for some scaling factor,  $\tau$ , in units of time, in order to write the system in dimensionless form.

2. **Modeling Traffic Flow.** Consider the initial value problem

$$\begin{cases} \frac{\partial u}{\partial t} + g'(u) \frac{\partial u}{\partial x} = 0; \\ u(x, 0) = f(x), \end{cases}$$

where  $g(u) = u(1 - u)$ , and the initial condition  $f$  is given by

$$f(x) = \begin{cases} 1, & \text{if } x < -1; \\ \frac{1}{2}(1 - x), & \text{if } -1 \leq x < 1; \\ 0, & \text{if } x \geq 1. \end{cases}$$

- Sketch the characteristic curves of the partial differential equation.
- Explain how the initial value problem can be solved in this case, and give a formula for  $u(x, t)$ .

3. **Age Structured Population Models.** Postulate a population density,  $n(a, t)$ , which also gives the age distribution for individuals in the population; so that, the number of individuals in the population between the ages  $a_1$  and  $a_2$  at time  $t$  is given by  $\int_{a_1}^{a_2} n(a, t) da$ .

- Explain why  $n(a, t)$  is given in units of population divided by units of time.
- Since  $a$  is a function of  $t$ , assuming that  $n$  is  $C^1$ , we can use Chain Rule to compute the rate of change of population density at time  $t$ ,  $\frac{dn}{dt}$ .

Explain why

$$\frac{dn}{dt} = \frac{\partial n}{\partial t} + \frac{\partial n}{\partial a}.$$

- Assume that death rate for individuals of age  $a$  in the population is proportional to the number of individuals at that age with constant of proportionality  $\mu(a)$ .

Use a conservation principle to derive the following partial differential equation

$$\frac{\partial n}{\partial t} + \frac{\partial n}{\partial a} = -\mu(a)n$$

Give the characteristic curves for the equation.

- Give solutions to the partial differential equation derived in the previous part assuming that the death rate is zero for all ages. Interpret your result.