

## Assignment #1

Due on Wednesday, January 28, 2015

Read Chapter 1 in the class lecture notes at <http://pages.pomona.edu/~ajr04747/>

Read Chapter 2, *Introduction: An Example from Epidemiology*, in the class lecture notes at <http://pages.pomona.edu/~ajr04747/>

**Background and Definitions.**

In Section 2.1 of the class lecture notes, we derived the Kermack–McKendrick SIR model for the spread of an infectious disease in a population,

$$\begin{cases} \frac{dS}{dt} = -\beta SI; \\ \frac{dI}{dt} = \beta SI - \gamma I; \\ \frac{dR}{dt} = \gamma I. \end{cases} \quad (1)$$

The quantity  $S(t)$  denotes the number of individuals in the population that are susceptible to getting the disease,  $I(t)$  is the number of individuals that have contracted the disease and can infect individuals from the susceptible class, and  $R(t)$  is the number of individuals in the population that have recovered and are immune to the disease. The positive parameters  $\beta$  and  $\gamma$  are called the infection rate and recovery rate, respectively.

Do the following problems

1. Give units for the parameters  $\beta$  and  $\gamma$  in the SIR system in (1).
2. Let  $N(t) = S(t) + I(t) + R(t)$  for all  $t$ . Use the equations in (1) to derive the differential equation

$$\frac{dN}{dt} = 0.$$

Deduce that  $N(t)$  must be a constant function.

3. Let  $S_o = S(t)$ , the initial number of susceptible individuals in the population and put

$$R_o = \frac{\beta S_o}{\gamma}. \quad (2)$$

Give the units for  $R_o$ .

The constant  $R_o$  is called the reproduction number.

4. Assume that at time  $t = 0$ , there are only one infectious individual in the population and no one in the population has acquired immunity. Let  $N$  denote the total number of individuals in the population.

- (a) Compute  $S_o$  in terms of  $N$ .
- (b) Give the reproduction number,  $R_o$ , in (2) in this situation.

5. Let  $R_o$  be as computed in Problem 4.

- (a) Assume that  $R_o > 1$ , and determine the sign of  $I'(0)$ . What do you conclude in this case? Explain the reasoning leading to your conclusion.
- (b) Assume that  $R_o < 1$ , and determine the sign of  $I'(0)$ . What do you conclude in this case? Explain the reasoning leading to your conclusion.