Review Problems for Exam 2

- 1. Consider a wheel of radius a which is rolling on the x-axis in the xy-plane. Suppose that the center of the wheel moves in the positive x-direction and a constant speed v_o . Let P denote a fixed point on the rim of the wheel.
 - (a) Give a path $\sigma(t) = (x(t), y(t))$ giving the position of the P at any time t, if P is initially at the point (0, 2a).
 - (b) Compute the velocity of P at any time t. When is the velocity of P horizontal? What is the speed of P at those times?
- 2. Let $f: \mathbb{R} \to \mathbb{R}$ denote a twice-differentiable real valued function and define

$$u(x,t) = f(x-ct)$$
 for all $(x,t) \in \mathbb{R}^2$,

where c is a real constant.

Show that

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}.$$

3. Let $f: \mathbb{R} \to \mathbb{R}$ denote a twice-differentiable real valued function and define

$$u(x,y) = f(r)$$
 where $r = \sqrt{x^2 + y^2}$ for all $(x,y) \in \mathbb{R}^2$.

Express the Laplacian of u, Δu , i.e., the divergence of the gradient of u, in terms of f', f'' and r.

- 4. Let f(x,y) = 4x 7y for all $(x,y) \in \mathbb{R}^2$, and $g(x,y) = 2x^2 + y^2$.
 - (a) Sketch the graph of the set $C = g^{-1}(1) = \{(x, y) \in \mathbb{R}^2 \mid g(x, y) = 1\}.$
 - (b) Show that at the points where f has an extremum on C, the gradient of f is parallel to the gradient of g.
 - (c) Find largest and the smallest value of f on C.
- 5. Let $C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1, y \ge 0\}$; i.e., C is the upper unit semi-circle. C can be parametrized by

$$\sigma(\tau) = (\tau, \sqrt{1 - \tau^2}) \text{ for } -1 \leqslant \tau \leqslant 1.$$

(a) Compute s(t), the arclength along C from (-1,0) to the point $\sigma(t)$, for $0 \le t \le 1$.

- (b) Compute s'(t) for -1 < t < t and sketch the graph of s as function of t.
- (c) Show that $\cos(\pi s(t)) = t$ for all $-1 \leq t \leq 1$, and deduce that

$$\sin(s(t)) = \sqrt{1 - t^2}$$
 for all $-1 \le t \le 1$.

6. Let R denote the open unit disc in \mathbb{R}^2 ; that is, $R = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$. Evaluate the integral

$$\int_{R} \ln(x^2 + y^2) \, \mathrm{d}x \, \mathrm{d}y$$

by first evaluating the integral

$$\int_{A_{\varepsilon}} \ln(x^2 + y^2) \, \mathrm{d}x \, \mathrm{d}y,$$

where A_{ε} is the annulus $\{(x, y) \in \mathbb{R}^2 \mid \varepsilon^2 < x^2 + y^2 < 1\}$, for $0 < \varepsilon < 1$, and then computing the limit at ε goes to 0.

7. Let A denote the annulus $\{(x,y) \in \mathbb{R}^2 \mid 1 < x^2 + y^2 < 4\}$, and evaluate $\int_A \frac{1}{x^2 + y^2} \, \mathrm{d}x \, \mathrm{d}y$.

8. Let
$$R = \{(x, y) \in \mathbb{R}^2 \mid 0 \le x \le y, x^2 + y^2 \le 1\}$$
, and evaluate $\int_R x^2 dx dy$.

- 9. Let *R* denote the region in the *xy*-plane bounded by the lines x + y = 1, x + y = 4, x y = -1 and x y = 1. Evaluate $\int_{R} (x + y)e^{x-y} dx dy$.
- 10. Evaluate $\int_{R} (x+y) dx dy$ where R is the rectangle in the xy-plane with vertices (1,0), (4,3), (3,4) and (0,1).
- 11. Evaluate $\int_{R} (x-y) dx dy$ where R is the square in the xy-plane with vertices (0,0), (2,-1), (3,1) and (1,2).
- 12. Let $\omega = 2x \, dx + y \, dy$ and $\eta = y \, dx x \, dy$ denote differential 1-forms. Compute each of the following $\omega \, d\eta$, $\eta \, d\omega$ and $d(\omega \eta)$.
- 13. Let C denote the unit circle traversed in the counterclockwise direction. Evaluate the line integral $\int_C x^3 dy y^3 dx$.
- 14. Let $F(x, y) = y \ \hat{i} x \ \hat{j}$ and R be the square in the xy-plane with vertices (0, 0), (2, -1), (3, 1) and (1, 2). Evaluate $\int_{\partial R} F \cdot n \, \mathrm{d}s$.