## Assignment #1

## Due on Wednesday, January 25, 2017

Read Chapter 1 in the class lecture notes at http://pages.pomona.edu/~ajr04747/

**Read** Chapter 2 on *Introduction to Modeling* in the class lecture notes at http://pages.pomona.edu/~ajr04747/

**Read** Section 1.1 on *Modeling via Differential Equations* in Blanchard, Devaney and Hall.

**Do** the following problems

1. The diagram in Figure 1 shows a simplification of the chemostat model discussed in Section 2.2 in the class lecture notes at http://pages.pomona.edu/~ajr04747/. The compartment in the diagram in Figure 1 represents a culture chamber con-

Co	
	$\begin{array}{c} N(t) \\ Q(t) \end{array}$

Figure 1: One–Compartment Model

taining N(t) bacteria and a quantity Q(t) of nutrient at time t. The quantities N and Q are assumed to be differentiable functions of t. Assume also that there is no flow of culture in or out of the chamber and that the culture in the chamber is kept well-stirred. In addition, assume that there is an initial amount of nutrient,  $Q_o$ , at an initial concentration of  $C_o$ , and that there are  $N_o$  bacteria at time t = 0. Postulate that the *per-capita* growth, K(c), is a function of the nutrient concentration,

$$c(t) = \frac{Q(t)}{V},$$

where V is the volume of the culture, which is assumed to be constant. Assuming that  $Y = 1/\alpha$  new cells are produced as s result of consumption of one unit of nutrient, apply conservations principles obtain a model for the evolution of N and Q in the chamber.

## Math 102. Rumbos

2. Combine the differential equations derived in Problem 1 to show that

$$\frac{d}{dt}[\alpha N + Q] = 0.$$

Deduce therefore that

$$\alpha N(t) + Q(t) = \alpha N_o + Q_o$$
, for all t.

3. Denote  $\alpha N_o + Q_o$  by  $A_o$  and use the result in Problems 2 to obtain the formula

$$c = \frac{A_o}{V} - \frac{\alpha}{V}N\tag{1}$$

for the concentration of nutrient.

- (a) Give an interpretation for the expression in (1).
- (b) Denote  $A_o/V$  by  $c_o$ . Explain why  $c_o$  is the nutrient concentration in the absence of bacteria.
- 4. Assume the constitutive equation K(c) = mc, where m is a positive constant of proportionality. Combine the results in Problems 1 and 3 to derive the differential equation

$$\frac{dN}{dt} = mN\left(c_o - \frac{\alpha}{V}N\right).$$
(2)

5. Set  $r = mc_o$  and  $L = \frac{c_o V}{\alpha}$ , and use (2) to derive a single equation describing the growth of bacteria in the chamber. This equation is known as the Logistic Equation.