

FINAL EXAM REVIEW

MATH 2110Q – Fall 2015

Professor Hohn

1. Find the values of x such that the vectors $\langle 3, 2, x \rangle$ and $\langle 2x, 4, x \rangle$ are orthogonal.
2. A constant force $\vec{F} = 3\hat{x} + 5\hat{y} + 10\hat{z}$ moves an object along the line segment from $(1, 0, 2)$ to $(5, 3, 8)$. Find the work done if the distance is measured in meters and force in newtons.
3. Let $f(x, y) = x^2 + xy$ on \mathbb{R}^2 , and let $F = \nabla f$. Let C be a curve in \mathbb{R}^2 starting at the point $(-1, -1)$ and ending at $(2, 3)$. Find $\int_C F \cdot d\vec{r}$.
4. Let $f(x, y) = x^2 + xy$ on \mathbb{R}^2 again. If C is the line segment starting at the origin $(0, 0)$ and ending at $(2, -1)$, find the line integral $\int_C f ds$.
5. Find the distance between the planes $3x + y - 4z = 2$ and $3x + y - 4z = 24$.
6. Find an equation of the plane through the line of intersection of the planes $x - z = 1$ and $y + 2z = 3$ and perpendicular to the plane $x + y - 2z = 1$.
7. Show that the planes $x + y - z = 1$ and $2x - 3y + 4z = 5$ are neither parallel nor perpendicular.
8. Find the length of the curve $\vec{r}(t) = \langle 2t^{3/2}, \cos(2t), \sin(2t) \rangle$, $0 \leq t \leq 1$.
9. Find a vector function that represents the curve of intersection of the cylinder $x^2 + y^2 = 16$ and the plane $x + z = 5$.
10. If $\vec{r}(t) = \langle t^2 + t \cos \pi t, \sin \pi t \rangle$, evaluate $\int_0^1 \vec{r}(t) dt$.
11. Let $f(x, y) = \sqrt{4 - x^2 - y^2}$.
 - (a) Find and sketch the domain of the function.
 - (b) Sketch 3 level curves of the surface described by the function.
 - (c) Find the first partial derivatives of the function.
 - (d) Find the tangent plane of the surface described by the function at the point $(1, 1, \sqrt{2})$
12. Find the points on the hyperboloid $x^2 + 4y^2 - z^2 = 4$ where the tangent plane is parallel to the plane $2x + 2y + z = 5$.
13. If $v = x^2 \sin y + ye^{xy}$, where $x = s + 2t$ and $y = st$, use the Chain Rule to find $\partial v / \partial s$ and $\partial v / \partial t$ when $s = 0$ and $t = 1$.
14. Let $f(x, y, z) = x^2y + x\sqrt{1+z}$.
 - (a) Find the directional derivative of f at the point $(1, 2, 3)$ in the direction $v = 2\hat{x} + \hat{y} - 2\hat{z}$.
 - (b) Find the maximum rate of change of f at the point $(1, 2, 3)$.
 - (c) In what direction does the maximum rate of change occur? Write your answer as a unit vector.

15. Find the absolute maximum and minimum values of $f(x, y) = e^{-x^2-y^2}(x^2 + 2y^2)$ on the disk $x^2 + y^2 \leq 4$.

16. A package in the shape of a rectangular box can be mailed by USPS if the sum of its length and girth (the perimeter of a cross section perpendicular to the length) is at most 108 in. Find the dimensions of the package with largest volume that can be mailed.

17. Find the local maximum and minimum values and saddle points of the function

$$f(x, y) = x^3 - 6xy + 8y^3.$$

18. Calculate the integral:

(a) $\int_0^1 \int_x^{e^x} 3xy^2 \, dy \, dx$

(b) $\int_0^1 \int_0^y \int_x^1 6xyz \, dz \, dx \, dy$

19. Describe/sketch the solid whose volume is given by the integral

$$\int_0^{\pi/2} \int_0^{\pi/2} \int_1^2 \rho^2 \sin \phi \, d\rho \, d\phi \, d\theta$$

and evaluate the integral.

20. Calculate the iterated integral by first reversing the order of integration.

$$\int_0^1 \int_x^1 \cos(y^2) \, dy \, dx$$

21. Find the volume of the solid that is bounded by the cylinder $x^2 + y^2 = 4$ and the planes $z = 0$ and $y + z = 3$.

22. Find the volume of the solid that is above the paraboloid $z = x^2 + y^2$ and below the half-cone $z = \sqrt{x^2 + y^2}$.

23. Convert the following integral into an integral with spherical coordinates.

$$\int_{-2}^2 \int_0^{\sqrt{4-y^2}} \int_{-\sqrt{4-x^2-y^2}}^{\sqrt{4-x^2-y^2}} y^2 \sqrt{x^2 + y^2 + z^2} \, dz \, dx \, dy$$

24. Evaluate the line integral

$$\int_C x \, ds$$

where C is the arc of the parabola $y = x^2$ from $(0, 0)$ to $(1, 1)$.

25. Compute the line integral

$$\int_C \vec{F} \cdot d\vec{r}$$

where $F(x, y) = \langle xy, x^2 \rangle$ and C is given by $\vec{r}(t) = \langle \sin t, 1 + t \rangle$, $0 \leq t \leq \pi$.

26. Find the work done by the force field

$$\vec{F}(x, y, z) = z\hat{x} + x\hat{y} + y\hat{z}$$

in moving a particle from the point $(3, 0, 0)$ to the point $(0, \pi/2, 3)$ along

- (a) a straight line
(b) the helix $x = 3 \cos t$, $y = t$, $z = 3 \sin t$.
27. Show that $F(x, y) = \langle (1+xy)e^{xy}, e^y + x^2e^{xy} \rangle$ is a conservative vector field. Then, find a function f such that $\vec{F} = \nabla f$.
28. Use Green's Theorem to evaluate

$$\int_C \sqrt{1+x^3} dx + 2xy dy$$

where C is the triangle with vertices $(0, 0)$, $(1, 0)$, and $(1, 3)$.

29. Let $\vec{F}(x, y, z) = \langle e^{-x} \sin y, e^{-y} \sin z, e^{-z} \sin x \rangle$.

- (a) Find the $\text{curl} \vec{F}$.
(b) Find the $\text{div} \vec{F}$.

30. If \vec{F} is a vector field defined on all of \mathbb{R}^3 whose component functions have continuous partial derivatives and $\text{curl} \vec{F} = 0$, then \vec{F} is a conservative vector field. Use this fact to show that $\vec{F}(x, y, z) = \langle e^y, xe^y + e^z, ye^z \rangle$ is conservative. Use the fact that \vec{F} is conservative to evaluate $\int_C \vec{F} \cdot d\vec{r}$ along the curve C where C is the line segment from $(0, 2, 0)$ to $(4, 0, 3)$.

Solutions (in no particular order):

$$\begin{aligned}
& f(x, y) = e^y + xe^{xy} + C; 2; 3; \langle -e^{-y} \cos z, -e^{-z} \cos x, -e^{-x} \cos y \rangle, -e^{-x} \sin y - e^{-y} \sin z - e^{-z} \sin x; \\
& -4, -2; 87; 8; \frac{2\sqrt{5}}{3}; 22/\sqrt{26}; x + y + z = 4; \frac{2}{27}(13^{3/2} - 8); \langle 4 \cos t, 4 \sin t, 5 - 4 \cos t \rangle, 0 \leq t \leq 2\pi; \\
& \frac{2}{9}e^3 - \frac{4}{45}; 1/4; \frac{7\pi}{6}; 1/2 \sin 1; 12\pi; \pi/6; \frac{64}{9}\pi; \frac{1}{12}(5\sqrt{5} - 1); \pi/4; \frac{1}{2}(3\pi - 9), \frac{-3\pi}{4}; \\
& \langle t^3/3, \frac{1}{\pi} \left(t \sin \pi t + \frac{\cos \pi t}{\pi} \right), \frac{-\cos \pi t}{\pi} \rangle; (2, 1/2, -1), (-2, -1/2, 1); 5, 0; 2e^{-1}, 0; 36, 18, 18; -1, (0, 0); \\
& \{(x, y) \mid x^2 + y^2 \leq 4\}; \frac{-y}{\sqrt{4 - x^2 - y^2}}; \frac{-x}{\sqrt{4 - x^2 - y^2}}; x + y + \sqrt{2}z = 4; 25/6; \sqrt{\frac{593}{16}}; \\
& \left\langle \frac{24}{\sqrt{593}}, \frac{4}{\sqrt{593}}, \frac{1}{\sqrt{593}} \right\rangle
\end{aligned}$$